



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by G. W. GREENWOOD, M. A. (Oxon), Lebanon, Ill.

Since P and D are extremities of conjugate diameters, their coördinates may be written $(a\cos\phi, b\sin\phi)$, $(a\sin\phi, -b\cos\phi)$ where a and b are the semi-axes. We obtain the required result at once by disregarding the sign and considering only the absolute magnitudes of the coördinates. Equations to AP and BD are

$$\frac{x - a\cos\phi}{a - a\cos\phi} = \frac{y - b\sin\phi}{-b\sin\phi}, \text{ and } \frac{x - a\sin\phi}{-a\sin\phi} = \frac{y + b\cos\phi}{b + b\cos\phi},$$

and the inclinations are therefore

$$-\frac{b\sin\phi}{a(1 - \cos\phi)} \text{ and } -\frac{b(1 + \cos\phi)}{a\sin\phi},$$

which are easily seen to be equal.

Also solved by G. B. M. Zerr.

CALCULUS.

187. Proposed by L. T. JACKSON, St. Louis, Mo.

Find the area of the ellipse

$$\begin{aligned} x &= a_1 + a_2 \cos\theta + a_3 \sin\theta, \\ y &= b_1 + b_2 \cos\theta + b_3 \sin\theta. \end{aligned}$$

I. Solution by H. B. LEONARD, B. S., Chicago, Ill.

Let $x' = x - a_1$, $y' = y - b_1$; then eliminating $\cos\theta$ and $\sin\theta$ in turn we obtain

$$b_2 x' - a_2 y' = \Delta \sin\theta, \quad b_3 x' - a_3 y' = -\Delta \cos\theta,$$

where $\Delta = b_2 a_3 - a_2 b_3$. The substitution $x'' = b_2 x' - a_2 y'$, $y'' = b_3 x' - a_3 y'$, which multiplies areas by Δ , transforms the ellipse into the circle $x'' = \Delta \sin\theta$, $y'' = -\Delta \cos\theta$; i. e. $x''^2 + y''^2 = \Delta^2$.

The area of circle being $\pi \Delta^2$, the area of ellipse is $\pi \Delta = \pi(b_2 a_3 - a_2 b_3)$.

II. Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Transform the origin to the center by writing x for $x - a_1$, y for $y - b_1$.

$$\therefore x = a_2 \cos\theta + a_3 \sin\theta, \quad y = b_2 \cos\theta + b_3 \sin\theta.$$

$$\begin{aligned} \text{Area} &= \int y dx = \int_0^{2\pi} (b_2 \cos\theta + b_3 \sin\theta)(a_3 \cos\theta - a_2 \sin\theta) d\theta \\ &= \int_0^{2\pi} (a_3 b_2 \cos^2\theta - a_2 b_3 \sin^2\theta - a_2 b_2 \sin\theta \cos\theta \pm a_3 b_3 \sin\theta \cos\theta) d\theta \\ &= \pi(a_3 b_2 - a_2 b_3). \end{aligned}$$

Also solved by F. P. Matz, and the Proposer.